

Comment on ‘The Newtonian force experienced by a point mass near a finite cylindrical source’

M Azreg-Aïnou

Başkent University, Department of Mathematics, Bağlıca Campus, Ankara, Turkey

Abstract. We compare the computation times and precisions of three expansions for the gravitational potential. The evaluation of the series by Selvaggi *et al* (2008 *Class. Quantum Grav.* **25** 015013) is time-consuming to be applied to the gravitational constant (G) or STEP experiments.

PACS numbers: 04.80.-y, 02.60.-x

We compare the computational efficiency of three published expansions for the gravitational potential (GP) of a finite, hollow, thick-walled cylinder with constant mass density ρ and reflection symmetry with respect to the plane $z = 0$. The expansions are those developed by Lockerbie *et al*, Schlamminger *et al* and Selvaggi *et al* [1, 2, 3].

If V denotes the GP/ $(2\pi G\rho)$ and $(2L, a, b)$ the length, inner and outer radii of the cylinder, Lockerbie *et al* formula, which is valid outside the sphere (S) of radius $\sqrt{b^2 + L^2}$ and center at the origin, writes in terms of Legendre polynomials P_{2n} as

$$V_{\text{Loc}}(r, z) = \sum_{n=0}^{\infty} \left\{ \sum_{p=0}^n \frac{(-1)^p (2n)! L^{2n-2p+1} (b^{2p+2} - a^{2p+2})}{2^{2p} p! (p+1)! (2n-2p+1)!} \right\} \frac{P_{2n}(z/\sqrt{r^2 + z^2})}{(r^2 + z^2)^{n+(1/2)}}, \quad (1)$$

The GP, $V_{\text{Sch}}(r, z)$, by Schlamminger *et al* formula is expressed in terms of the GP along the axis of symmetry $V(0, z)$. If we introduce $f(x) = x + \sqrt{b^2 + x^2}$ and $g(x) = x\sqrt{b^2 + x^2}$, then $V(0, z)$ & $V_{\text{Sch}}(r, z)$ are evaluated by

$$2V(0, z) = b^2 \{ \ln[f(z+L)] - \ln[f(z-L)] \} + g(z+L) - g(z-L) - (a \rightarrow b), \quad (2)$$

$$V_{\text{Sch}}(r, z) = V(0, z) + \sum_{n=1}^{\infty} \frac{(-1)^n V^{(2n)}(0, z)}{2^{2n} (n!)^2} r^{2n}, \quad (\text{Converges for small } r). \quad (3)$$

where $(a \rightarrow b)$ substitutes a for b in the first four terms of (2).

In terms of the hypergeometric function ${}_2F_1$ [4], Selvaggi *et al* formula reads

$$G(r, u, n, a, b) = b^{2n+2} {}_2F_1 \left(n+1, 2n+(1/2); n+2; -b^2/(u^2 + r^2) \right) - (a \rightarrow b),$$

$$V_{\text{Sel}}(r, z) = \frac{1}{2} \sum_{n=0}^{\infty} \frac{(4n-1)!!}{(n+1)(n!)^2} \left(\frac{r}{2} \right)^{2n} \int_{z-L}^{z+L} \frac{G(r, u, n, a, b)}{(u^2 + r^2)^{2n+(1/2)}} du. \quad (4)$$

We consider the case where $a = 1/2$, $b = 1$ & $L = 1$, in SI units. We restrict ourselves to the central plane $z = 0$ and compare the three formulae (1), (3) & (4) in their respective regions of convergence. We choose three points: $P_1(0.2, 0)$ near

the z -axis, $P_2(0.4, 0)$ near the inner surface of the cylinder and $P_3(2, 0)$ outside the cylinder. If $V(P, n+1)$ denotes the potential at the point P with $n+1$ being the number of terms used (here, n represents the index of summation which has been given the same symbol in all formulae (1), (3) & (4)), our results follow where the exact decimal places are underlined

$$V_{\text{Sel}}(P_1, 4) \simeq 0.\underline{820253}; V_{\text{Sel}}(P_1, 7) \simeq 0.\underline{820382}; V_{\text{Sch}}(P_1, 3) \simeq 0.\underline{820385}, \quad (5)$$

$$V_{\text{Sel}}(P_2, 5) \simeq 0.\underline{825570}; V_{\text{Sel}}(P_2, 7) \simeq 0.\underline{828583}; V_{\text{Sch}}(P_2, 5) \simeq 0.\underline{831536}, \quad (6)$$

$$V_{\text{Sel}}(P_3, 7) \simeq 0.\underline{371846}; V_{\text{Loc}}(P_3, 7) \simeq 0.\underline{371958}. \quad (7)$$

$V_{\text{Sch}}(P_3)$ blows up and so do $V_{\text{Loc}}(P_1)$ & $V_{\text{Loc}}(P_2)$. The number of exact decimal places in (5) to (7) has been identified upon comparing $V_{\text{Sch}}(P_1, 3)$, $V_{\text{Sch}}(P_2, 5)$ & $V_{\text{Loc}}(P_3, 7)$ to more accurate results, which have been obtained at the same points using large values of $n+1$ (see last paragraph). The relative errors at P are $E_{\text{Sel-Sch}}(P_1, 4) \simeq 1.61 \times 10^{-4}$, $E_{\text{Sel-Sch}}(P_2, 7) \simeq 3.55 \times 10^{-3}$ & $E_{\text{Sel-Loc}}(P_3, 7) \simeq 3.01 \times 10^{-4}$ with $E_{\text{Sel-Sch}}(P, n+1) = |V_{\text{Sel}}(P, n+1) - V_{\text{Sch}}(P, n+1)|/V_{\text{Sch}}(P, n+1)$, etc.

Near the plane $z = 0$, the expansions (1) & (3) perform better than (4) (Eqs (5) to (7)) and are easily handled with reasonable evaluation times. The evaluation of each term in (4) involves a non-trivial integration which is time-consuming. In contrast, (1) & (3) are evaluated via the use of Legendre polynomials, which are built-in functions in most computer algebra systems, or by differentiation. For instance, the evaluation of $V_{\text{Sel}}(P_2, 7)$, by numerical integration, lasted more than 29000 times that of $V_{\text{Sch}}(P_2, 7)$ (if we re-derive (2) using computer-algebra, this ratio reduces to 370) and the evaluation of $V_{\text{Sel}}(P_3, 7)$ lasted more than 46000 times that of $V_{\text{Loc}}(P_3, 7)$.

In the plane $z = 0$ each term of (4) ($n \geq 1$ up to at least 5) reaches a maximum value for $a < r_0 < b$ and decays slowly near and off the axis resulting in a slow convergence of (4) around r_0 . Off the plane $z = 0$, the convergence of (4) improves around the axis due to an off-axis shift of r_0 and a drop in the maximum resulting in a rapid decay of its terms ($n \geq 1$) there. At the point $P_4(0.4, 2)$, we found $V_{\text{Sel}}(P_4, 7) = 0.\underline{36482771}$, $V_{\text{Sch}}(P_4, 7) = 0.\underline{36482771}$ & $V_{\text{Loc}}(P_4, 7) = 0.\underline{36482807}$. The evaluation of $V_{\text{Sel}}(P_4, 7)$ lasted 8500 times that of $V_{\text{Sch}}(P_4, 7)$ and 52000 times that of $V_{\text{Loc}}(P_4, 7)$.

The flexibility in using (1) or (3) allowed as to evaluate the GP using up to forty-one terms and to check over the stability of the results shown in (5) to (7) for V_{Sch} & V_{Loc} . Moreover, the GP at each of the points P_1 to P_4 has been reobtained by numerically integrating the volume integral of the potential element $dv'/(2\pi|\mathbf{r} - \mathbf{r}'|)$.

References

- [1] Lockerbie N A, Veryaskin and Xu X 1993 *Class. Quantum Grav.* **10** 2419-2430
- [2] Schlaminger St, Holzschuh E, Kündig W, Nolting F, Pixley R E, Schurr J and Straumann U 2006 *Phy. Rev. D* **74** 082001 (25 pp)
- [3] Selvaggi J P, Sheppard S and Chari M V K 2008 *Class. Quantum Grav.* **25** 015013 (15 pp)
- [4] <http://functions.wolfram.com/PDF/Hypergeometric2F1.pdf>